Delivering in the Time of COVID-19: A Population-Based Study of Childbirths in the U.S.

Background

Since December 2019, COVID-19 (new coronavirus pneumonia) has spread widely, posing a great threat to public health worldwide. On March 11, 2020, the World Health Organization announced that SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus Type 2) was spreading globally, known as the COVID-19 outbreak [1]. The epidemic has caused unprecedented damage to health care system in many countries and regions. The morbidity and mortality associated with SARS-CoV-2 infection has placed enormous strain on healthcare systems and has caused significant disruption to the delivery of non-COVID-19 healthcare services [2]. The suddenness and severity of the outbreak have adversely affected a variety of health care facilities and services, forcing all healthcare professionals and patients to reconsider how and how often healthcare is provided [3].

These disruptions to health services have led to serious collateral damage, disproportionately affecting vulnerable populations, especially pregnant women and newborns, despite of various emerging clinical guidelines for obstetric care [4, 5]. Across the world, multiple studies have shown that neonatal mortality and stillbirth rates have risen sharply, and birth rates have declined during the COVID-19 pandemic, regardless of maternal COVID-19 infection [6]. Based on data analysis of multiple regions before and after the epidemic [7, 8], studies suggest that quality of childbirth care services are influenced by human factors of healthcare professionals--- staff burnout, culture, teamwork [9], persisting high pressure at work [10-12]---as well as birth companions from family and other social connections [13].

Objectives

We aim to investigate the impacts of the COVID-19 pandemic on childbirth outcomes, maternal and newborn healthcare in the U.S. By identifying changes in childbirths and associated health services before and after the onset of COVID-19, we provide valuable managerial insights for healthcare administration and professionals to provide high-quality health care for mothers and infants. The problems exposed by this global epidemic and corresponding improvements will contribute to precautious planning in possible national or global health crises in the future.

TEXAS

UNIVERSITY

Dr. Emily Zhu Fainman

Department of CIS & Analytics, McCoy College of Business, Texas State University



Contact

Dr. Emily Zhu Fainman Assistant Professor



mathematics, including analytics, statistics, operations management and economics. Closely collaborating with physicians, health administration, and researchers, she is working on several projects related to maternity care, oncology, Emergency Departments, electric vehicles and shared transport. Dr. Zhu received her Ph.D. in Operations Management from McGill University, Canada in 2018; she has a M.Sc. in Mathematical and Computational Finance from University of Oxford, UK, and a B.Sc. in Mathematics from Nanjing University, China.

TEXAS STATE

UNIVERSITY

References

[1] WHO. WHO Director-General's opening remarks at the media briefing on March 2020[R]. Geneva:WHO,2020.

[2] Tanne J. H., Hayasaki E., Zastrow M., et al. Covid-19: how doctors and healthcare systems are tackling coronavirus worldwide. BMJ. 2020:368.

[3] Semaan A., Audet C., Huysmans E., et al. Voices from the frontline: findings from a thematic analysis of a rapid online global survey of maternal and newborn health professionals facing the COVID-19 pandemic. BMJ Glob Health. 2020,5(6).

[4] Boelig R. C., Saccone G., Bellussi F., et al. MFM guidance for COVID-19. Am J *Obstet Gynecol MFM*. 2020,2(2):100-106.

[5] Khalil A., von Dadelszen P., Kalafat E., et al. Change in obstetric attendance and activities during the COVID-19 pandemic. Lancet Infect Dis. 2021,21(5):e115.

[6] Meyer R., Bart Y., Tsur A., et al. A marked decrease in preterm deliveries during the coronavirus disease 2019 pandemic. Am J Obstet Gynecol. 2021,224(2):234-7.

[7] Amouzou A., Maiga A., Faye C. M., et al. Health service utilisation during the COVID-19 pandemic in sub-Saharan Africa in 2020: a multicountry empirical assessment with a focus on maternal, newborn and child health services. BMJ Glob Health. 2022,7(5).

[8] Aranda Z., Binde T., Tashman K., et al. Disruptions in maternal health service use during the COVID-19 pandemic in 2020: experiences from 37 health facilities in lowincome and middle-income countries. *BMJ Glob Health*. 2022,7(1).

[9] Tawfik D. S., Scheid A., Profit J., et al. Evidence relating health care provider burnout and quality of care: a systematic review and meta-analysis. Ann Intern Med. 2019,171(8):555-567.

[10] Hartzband P. and Groopman J. Physician burnout, interrupted. N Engl J Med. 2020,382(26):2485-2487.

[11] Krystal J. H. Responding to the hidden pandemic for healthcare workers: stress. Nat Med. 2020,26(5):639-639.

[12] Schmitt N., Mattern E., Cignacco E., et al. Effects of the Covid-19 pandemic on maternity staff in 2020 - a scoping review. BMC Health Serv Res. 2021,21(1):1364. [13] WHO. WHO recommendations: intrapartum care for a positive childbirth

experience: web annex; evidence base[R]. Geneva:WHO,2018.

research focuses on Dr. Zhu's effectiveness improving and efficiency in healthcare and service contributes to She systems. strategic, operational and technical policy-making applying by interdisciplinary approaches,

MEMBER THE TEXAS STATE UNIVERSITY SYSTEM